



細胞性粘菌由来の新規低分子化合物Ppc-1による体重減少

メタデータ	言語: English 出版者: 公開日: 2015-10-22 キーワード (Ja): キーワード (En): 作成者: 鈴木, 俊幸 メールアドレス: 所属:
URL	https://fmu.repo.nii.ac.jp/records/2000140

学位論文

Weight Loss by Ppc-1, a Novel Small Molecule Mitochondrial Uncoupler
Derived from Slime Mold
(細胞性粘菌由来の新規低分子化合物 Ppc-1 による体重減少)

医学部附属放射性同位元素研究施設

鈴木俊幸

Weight Loss by Ppc-1, a Novel Small Molecule Mitochondrial Uncoupler
Derived from Slime Mold
(細胞性粘菌由来の新規低分子化合物 Ppc-1 による体重減少)

医学部附属放射性同位元素研究施設

鈴木俊幸

概要

ミトコンドリアの主な機能は電子伝達系における酸化リン酸化による ATP 産生であるが、脂質や糖質の代謝、細胞周期やアポトーシスなどにおいても重要な役割を担っている。これらの機能を制御する新規化合物は、新しい抗がん剤などの開発に繋がる可能性があり、また、分子メカニズム解析のツールとして有用である。

細胞性粘菌は土壌微生物で、薬理学的作用を持つ多くの化合物を産生することから、医学研究に用いられる化合物の重要な資源の一つである。細胞性粘菌 *Dictyostelium discoideum* (キイロタマホコリカビ) 由来の分化因子 differentiation inducing factor-1(DIF-1)とその代謝産物である DIF-3 は、細胞増殖抑制作用があり抗腫瘍作用が期待されている。近年、DIF-3 とそのアナログがミトコンドリアへも作用していることが報告された。

本研究では、細胞性粘菌および微生物由来の低分子化合物について、ミトコンドリアにおける酸素消費率へおよぼす効果を指標にスクリーニングを行い、新規ミトコンドリア標的化合物を探索した。その結果、細胞性粘菌 *Polysphondylium pseudo-candidum* (リュウキュウカビモドキ) 由来の Ppc-1 を、酸素消費率を亢進するが ATP 産生に顕著な影響を与えない化合物として同定した。さらに、Ppc-1 の作用を細胞および個体レベルで解析した。

細胞性粘菌由来の 9 種を含む 32 種の天然低分子化合物について、マウス肝臓から調整したミトコンドリア画分の酸素消費率を指標にスクリーニングを行った。これらのうち、6 種の化合物が ADP 非存在下 (state 4) での酸素消費率を増加させたが、5 つは ADP が存在下 (state 3) の酸素消費率を阻害した。Ppc-1 のみが state 3 を阻害しなかった。また、ATP の合成および分解に影響を与えなかった。Ppc-1 による酸素消費率の増加は、シトクロムオキシダーゼ阻害剤であるシアン化カリウムによって抑制されたが、ATP 合成酵素阻害剤であるオリゴマイシンは効果がなかった。ATP 合成酵素の ADP に対する K_m は、Ppc-1 の濃度を変えても 10~15 μM となり顕著な変化は認められなかった。これらの結果から、Ppc-1 はミトコンドリアにおける酸化リン酸化の脱共役剤であることが示唆された。

脱共役剤は、ミトコンドリア内膜を介したプロトン勾配を減少し ATP 産生の効率を低下することから、動物への投与により体重減少をひき起こすことが期待される。そこで Ppc-1 を週に一度腹腔内に 8 週間投与した。投与は、週一度のペースで 0 (コントロール)、0.16 (グループ A)、0.4 (同 B)、4 (同 C)、10 mg/週/kg 体重 (同 D) の容量で行い、各個体の体重を 1 日おきに消灯前に測定した。その結果、最終投与後の体重が、グループ B はコントロールに比べ 7~10% 低下した。グループ C については有意差が認められたものの、グループ B の場合よりも差は小さく、A および D では有意差はなかった。投与マウスの肝臓と腎臓の組織像は正常で、脂肪蓄積や細胞傷害は認められなかった。また、血清中アラニンアミノトランスフェラーゼ、血液尿素窒素、アディポネクチンなどの値も正常であった。飼育期間中の摂食量に有意差はなく、異常行動、腫瘍形成は観察されなかった。

Ppc-1 は 10 μM 以上の濃度で各種培養細胞の増殖を抑制することから、Ppc-1 (0.8mg/kg 体重) 投与後の血清および各種組織の Ppc-1 濃度を経時的に測定した。その結果、各組織の Ppc-1 濃度は投与後 30 分で最高値 ($\sim 0.1 \mu\text{M}$) を示し、投与後 24-48 時間後に検出限界以下

となった。これらの結果より、体重増加抑制が細胞増殖の抑制によるものではないことが示された。Ppc-1 投与マウスでは、血清中の遊離脂肪酸値および中性脂肪値が増加しており、X線CTによる内臓脂肪と皮下脂肪もコントロールに比べ少ない傾向を示した。脂肪細胞に分化させた3T3-L1細胞へPpc-1投与することにより、培地中の遊離脂肪酸が上昇したことから、Ppc-1投与マウスで観察された血清遊離脂肪酸の増加は、Ppc-1が脂肪細胞に作用して脂肪酸遊離を促進したことによるものと推測された。

以上の結果より、Ppc-1は新規脱共役物質で、ミトコンドリアのエネルギー代謝系の解析に有用であり、また、新しい肥満治療薬のリード化合物となる可能性が示唆された。

RESEARCH ARTICLE

Weight Loss by Ppc-1, a Novel Small Molecule Mitochondrial Uncoupler Derived from Slime Mold

Toshiyuki Suzuki¹, Haruhisa Kikuchi², Masato Ogura¹, Miwako K. Homma¹, Yoshiteru Oshima², Yoshimi Homma^{1*}

¹ Fukushima Medical University School of Medicine, Fukushima, 960–1295, Japan, ² Graduate School of Pharmaceutical Sciences, Tohoku University, Sendai, 980–8678, Japan

* yoshihom@fmu.ac.jp



CrossMark
click for updates

OPEN ACCESS

Citation: Suzuki T, Kikuchi H, Ogura M, Homma MK, Oshima Y, Homma Y (2015) Weight Loss by Ppc-1, a Novel Small Molecule Mitochondrial Uncoupler Derived from Slime Mold. PLoS ONE 10(2): e0117088. doi:10.1371/journal.pone.0117088

Academic Editor: Joseph Najbauer, University of Pécs Medical School, HUNGARY

Received: June 27, 2014

Accepted: December 19, 2014

Published: February 10, 2015

Copyright: © 2015 Suzuki et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Funding: This work was supported in part by a Grant-in-aid for Scientific Research from the Japan Society for the Promotion of Science (to YO: No. 24590110, and to HK: No. 23710247) and a grant from Fukushima Medical University (to YH: No. H2401). The funders had no role in study design, data collection and analysis, the decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Mitochondria play a key role in diverse processes including ATP synthesis and apoptosis. Mitochondrial function can be studied using inhibitors of respiration, and new agents are valuable for discovering novel mechanisms involved in mitochondrial regulation. Here, we screened small molecules derived from slime molds and other microorganisms for their effects on mitochondrial oxygen consumption. We identified Ppc-1 as a novel molecule which stimulates oxygen consumption without adverse effects on ATP production. The kinetic behavior of Ppc-1 suggests its function as a mitochondrial uncoupler. Serial administration of Ppc-1 into mice suppressed weight gain with no abnormal effects on liver or kidney tissues, and no evidence of tumor formation. Serum fatty acid levels were significantly elevated in mice treated with Ppc-1, while body fat content remained low. After a single administration, Ppc-1 distributes into various tissues of individual animals at low levels. Ppc-1 stimulates adipocytes in culture to release fatty acids, which might explain the elevated serum fatty acids in Ppc-1-treated mice. The results suggest that Ppc-1 is a unique mitochondrial regulator which will be a valuable tool for mitochondrial research as well as the development of new drugs to treat obesity.

Introduction

Mitochondria are organelles comprised of an outer and inner membrane, enclosing a matrix space. Within the inner membrane is a series of redox enzymes comprising the respiratory chain, which reduces oxygen to water using electrons generated by dehydrogenases employing intermediary metabolite substrates. The respiratory enzyme complexes are redox-driven proton pumps, which translocate protons from the matrix space into the inner mitochondrial compartment, thus establishing an electrochemical potential gradient across the inner mitochondrial membrane. The electrochemical gradient is utilized by the reversible proton pumping function of ATP synthase to synthesize ATP from ADP and inorganic phosphate. Various small compounds have been used to study mitochondrial regulation [1–3]. Rotenone [4] and

antimycin A [5] are well-known inhibitors of respiratory chain complex I and III, respectively, while oligomycin A is a specific inhibitor of ATP synthase. In addition, uncoupling agents such as 2,4-dinitrophenol (DNP) [6] and carbonyl cyanide m-chlorophenyl hydrazone (CCCP) [7] are useful probes for understanding the mechanisms involved in electron transfer and ATP production. DNP and CCCP are well-characterized mitochondrial uncoupling agents, and act as a protonophore to shuttle protons across the mitochondrial inner membrane, leading to dissipation of the mitochondrial proton gradient and conversion of the energy derived from mitochondrial substrate oxidation into heat.

Cellular slime molds are soil microorganisms that produce many pharmacologically active compounds, and are an important source of lead compounds for medical research. The slime mold *Dictyostelium discoideum* forms a fruiting body consisting of spores and a multicellular stalk at the end of its life cycle, in response to differentiation inducing factor-1 (DIF-1), a putative morphogen which induces stalk cell differentiation [8]. The first metabolite produced during DIF-1 degradation is DIF-3, which unlike DIF-1 has virtually no activity in the induction of stalk cell differentiation [9]. Nevertheless, both DIF-1 and DIF-3 possess anti-tumor activity, suppressing mammalian cell proliferation and, in some cases, inducing or promoting the differentiation of de-differentiated tumor cells *in vitro* [10–16]. Recently, we showed that DIF-3 and its analogues induce morphological changes and dysfunction of mitochondria, suggesting that DIF-like molecules suppress cell proliferation, at least in part, by interfering with mitochondrial activity [17,18]. In this study, we screened natural low-molecular-weight compounds derived from *D. discoideum* and other microorganisms to identify novel mitochondrial modulators. We report that the secondary metabolite, Ppc-1, is a unique compound which enhances mitochondrial oxygen consumption and induces weight loss in mice. Thus, Ppc-1 is novel and potentially valuable small molecule reagent for mitochondrial research.

Results

Oxygen consumption of natural compounds

We screened for novel modulators of mitochondrial function by measuring oxygen consumption rates in mitochondria-enriched fractions prepared from mouse liver. We tested 32 natural small molecule compounds, including 9 derived from slime molds (Fig. 1A) [8,9,19–24]. Assays were carried out in buffer containing 100 μ M palmitoyl-carnitine as a substrate for oxidation, and monitoring oxygen levels using an oxygen electrode. After preincubating the mitochondrial-enriched fraction in an assay buffer at 30°C for 30 s, each test compound was added to a final concentration of 20 μ M, followed by addition of ADP to a final concentration of 400 μ M, in order to induce 'state 3'. Four of the nine compounds derived from slime molds enhanced the oxygen consumption rate in the absence of ADP, including #11, #12, #27, and #28 (Ppc-1). We also examined 23 natural compounds derived from other microorganisms and plants, and found 2 more compounds with oxygen consumption-stimulating activity. Of the 6 selected compounds, five significantly inhibited the rate of 'state 3' (Fig. 1B). Only Ppc-1 lacked inhibition of the rate of 'state 3', therefore, we selected this compound for further analysis.

Ppc-1 enhanced oxygen consumption rate in a dose-dependent manner (Fig. 2A). The effects of oligomycin A and KCN were used to confirm the effects of Ppc-1 on respiration; while the cytochrome c oxidase inhibitor, KCN, completely suppressed the effect of Ppc-1 on oxygen consumption rate, the ATP synthase inhibitor, oligomycin A, had no effect (Fig. 2B). Whereas Ppc-1 concentrations above 10 μ M significantly enhanced oxygen consumption rate (Fig. 2C), no significant inhibition of 'state 3' was observed, even at such high concentration (Fig. 2D). The respiratory control ratio (RCR) for ADP suggests that Ppc-1 acts as a mitochondrial




Figure 1. Screening of uncoupling agents. (A) Structures of small molecule compounds identified in slime molds. (B) Effects of compounds (20 μ M each) on oxygen consumption were measured in the absence or presence (state 3) of ADP using isolated mitochondria. Four compounds were found to enhance oxygen consumption, suggesting uncoupling activity. Only Ppc-1 exhibited no inhibitory effect on 'state 3'.

doi:10.1371/journal.pone.0117088.g001

uncoupler (Fig. 2E). The apparent K_m for ADP was 10–15 μ M in the presence of varying concentrations of Ppc-1 (Fig. 2F), which were consistent with previous reports [25,26]. This confirms that Ppc-1 does not affect the function of ATP synthase.

Ppc-1 suppresses weight gain in animals

Uncoupling agents reduce the mitochondrial proton gradient, decreasing the efficiency of ATP production. It is well known that chronic administration of uncoupling agents in animals causes significant weight loss. Therefore, we examined the effect of Ppc-1 administration on weight and fat content in mice. ICR mice were divided into 5 groups (3–9 animals in each group), and Ppc-1 dissolved in phosphate buffered saline (PBS) was injected into the peritoneal cavity of each mouse once a week for 8 weeks. The Ppc-1 doses used were 0 (control), 0.16 (group A), 0.8 (group B), 4 (group C), and 10 mg/week/kg (group D). Since mice feed in the dark, body weight was measured every 2 days just before the lights were turned off. The most significant suppression of weight gain was observed in the group B mice (Fig. 3A), 3 weeks




Figure 2. Oxygen consumption induced by Ppc-1. (A) Enhanced oxygen consumption was induced by Ppc-1 in a dose-dependent manner. Mitochondria-enriched fraction (0.1 mg protein) in assay buffer was pre-incubated at 30°C for 0.5 min, and the indicated amounts of Ppc-1 were added to the mixture at time point 0 min. Recording of oxygen level was started at 0.5 min after the addition of the compound, and then an aliquot of ADP was added to a final concentration of 400 μM to induce 'state 3'. (B) Effects of oligomycin A (0.75 μM) and KCN (1 mM) on the oxygen consumption were examined. (C, D) The oxygen consumption rate in the absence (C) or presence (D) of ADP was calculated. Ppc-1 showed a significant increase in oxygen consumption at concentrations above 10 μM, but no effect on 'state 3' at any dose tested. ** $p < 0.01$. (E) The RCR for ADP was calculated, confirming that Ppc-1 acts as a mitochondrial uncoupler. (F) Effect of Ppc-1 on the oxygen consumption rate was examined in assay buffer containing various concentrations of ADP. A representative reciprocal plot is shown, and the K_m values for ADP are inserted.

doi:10.1371/journal.pone.0117088.g002

after the start of Ppc-1 administration, and this suppression was maintained throughout the experiment. Mean weights for the control and group B mice were respectively 32.3 g and 30.1 g after 5 weeks, and respectively 33.2 g and 30.6 g after 7 weeks, indicating 7~10% weight loss in Ppc-1 treated animals. Significant weight loss with Ppc-1 was also observed in the group C mice (Fig. 3B), although the effect was less pronounced. The weight differences between treated and the control animals were sustained, even after Ppc-1 administration was discontinued. On the other hand, weight differences were not significant between the control and group A or group D mice (S1 Fig.).




Figure 3. Suppression of weight gain by Ppc-1. Ppc-1 was injected once a week into the peritoneal cavity of mice for 8 weeks. Doses of Ppc-1 for groups B and C were 0.8 (A) and 4 (B) mg/week/kg, respectively. Results are expressed using box plots. ** $p < 0.01$, * $p < 0.05$.

doi:10.1371/journal.pone.0117088.g003

Increase in serum fatty acid levels by Ppc-1 treatment

We next compared components of serum from the control and group B mice. Three mice from each group were selected at random, and serum was collected from each animal on the 5th day after the final administration of Ppc-1. As shown in Table 1 and S1 Table, triacylglycerol and free fatty acids were significantly increased in Ppc-1-treated animals, suggesting enhanced lipid

Table 1. Profiling of Serum Factors.

	Triglyceride (mg/l)		Free fatty acid (μ EQ/l)		Adiponectin (mg/l)		BUN (mg/l)		AST (IU/l)		ALT (IU/l)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Control	733	61	508	117	17.1	3.4	229	71	54.7	15.4	6.3	3.5
Group A	740	85	582	83	17.0	2.5	204	30	72.3	17.1	6.4	3.6
Group B	943*	102	666*	84	16.4	4.0	229	18	48.0	17.8	3.7	0.6
Group C	653	253	893*	187	18.1	3.9	230	24	57.0	6.1	4.3	0.6
Group D	718	299	990	444	14.6	1.1	260	51	83.5	44.5	12.0	7.1

*: $p < 0.05$.

doi:10.1371/journal.pone.0117088.t001

metabolism in response to the compound. Analysis of serum by mass spectrometry (MS) also revealed enhancement of specific fatty acids in response to Ppc-1, particularly stearic acid (18:0) and oleic acid (18:1) (Fig. 4A). These results were confirmed by computed tomography (CT) scanning (Fig. 4B and S2 Fig.) where the subcutaneous and visceral fat levels in the group B mice were respectively $7.2 \pm 2.8\%$ and $7.9 \pm 3.6\%$, lower than the values measured in control mice of $9.9 \pm 1.4\%$ and $11.6 \pm 5.1\%$ (mean \pm SD, $n = 3$). Furthermore, in contrast to the control animals, group B mice showed no ectopic fat accumulation.

After serum collection, three mice from each group were euthanized and examined for tissue damage to liver and kidney. Tissue images are shown in Fig. 5A. Both liver and kidney images appeared normal, and no lesional changes were detected, including ectopic accumulation of fat. The remaining control and group B mice were housed under normal conditions and examined periodically until they died of natural causes. The levels of daily food intake were comparable among the control and Ppc-1 treatment groups (Fig. 5B). No abnormal behavior, evidence of tumor formation, or differences in life span was observed between the animals in the control and the four Ppc-1 treatment groups.

Tissue distribution of Ppc-1

Although Ppc-1 functions as an uncoupler and enhances oxygen consumption in mitochondria (Fig. 2C), no increase in body temperature was detected in mice treated with the reagent (Fig. 5C). However, Ppc-1 caused growth inhibition when added to mammalian cells at concentrations above $10 \mu\text{M}$ [19] (S3 Fig.). Therefore, we determined Ppc-1 concentrations within various tissues in mice treated with a single injection of Ppc-1 (0.8 mg/kg), collecting liver, kidney, visceral fat, and serum from the animals at different times after injection. Lipids were extracted from tissues with chloroform/methanol, and analyzed by LC-MS. The lipid fraction from RPE cells treated with $10 \mu\text{M}$ Ppc-1 for 12 h was also extracted and analyzed by LC-MS. A mass spectrum of Ppc-1 in RPE cells is shown in Fig. 6A; the major peak at m/z 356.19 was identified as intact Ppc-1 $[\text{M}+\text{H}]^+$, and other peaks (e.g. m/z 288.12) were recognized as derivatives of Ppc-1. Ppc-1 concentrations were low in the sera and tissues derived from mice administered the compound. At 1 h after animal injection, Ppc-1 was detected in serum and fat tissue, at concentrations (68 and 2.6×10^5 peak area/mg, respectively) which were less than 1% of that seen in cultured cells treated with $10 \mu\text{M}$ Ppc-1 (0.82×10^9 peak area/mg) (Fig. 6B). The signal intensity of Ppc-1 in tissues declined gradually to basal levels, 24 to 48 h after injection. The Ppc-1 concentrations in liver, kidney, and subcutaneous fat were extremely low throughout the time points examined.

Ppc-1 induces fatty acid release from adipocytes

In an effort to understand why serum fatty acid levels are elevated in Ppc-1-treated mice (Table 1), the effects of the compound on adipocytes were examined using differentiated 3T3-L1 cells. Treatment of 3T3-L1 adipocytes with $10 \mu\text{M}$ Ppc-1 for 2–3 days led to significantly increased levels of free fatty acids in the culture medium, suggesting that Ppc-1 stimulates fatty acid release from adipocytes (Fig. 7). Therefore, it is possible that in animals, Ppc-1 entering the bloodstream from the peritoneal cavity stimulates adipocytes directly, promoting their release of fatty acids.

Discussion

Identification of Ppc-1 as a novel mitochondrial uncoupler

Small molecules that inhibit mitochondrial respiration have been indispensable for investigating mitochondrial function and regulation, and new modulators offer possibilities for uncovering




Figure 4. Fat analysis of Ppc-1-treated mice. (A) Free fatty acids in the serum of mice administered Ppc-1 at a dose of 0.8 mg/week/kg for 8 weeks were analyzed by HPLC-MS/MS system A. (B) Whole body fat was examined by a CT scanner, and representative vertical section images are shown. Fat contents were quantified from the cross section images, and the results are expressed using box plots. The position analyzed is denoted by an arrow. * $p < 0.05$.

doi:10.1371/journal.pone.0117088.g004




Figure 5. Examination of Ppc-1-treated mice. (A) Cross sections were prepared of fixed kidney and liver tissues from mice in the control and Ppc-1 treated (0.8 mg/week/kg) groups and stained with hematoxylin-eosin. Representative images are shown; no significant differences were detected between the two groups. (B) Daily food intake in 4 groups was recorded throughout the experiments. (C) Body surface temperature was monitored with an infrared thermometer following a single injection of Ppc-1 (0.8 mg/kg).

doi:10.1371/journal.pone.0117088.g005

mechanisms which are as yet undiscovered. In our screen of 32 natural compounds, Ppc-1 was especially interesting as a compound with oxygen consumption-stimulating activity but without inhibitory effect on 'state 3'. 'State 3' is induced when both respiratory substrates and ADP are supplied to mitochondria. The electron transfer system and ATP synthase are strictly regulated, through molecular interactions and intra-mitochondrial signaling systems which under various conditions can be labile, reducing ATP synthesis and producing excess reactive oxygen species [27–30]. Ppc-1 at >98% purity enhances oxygen consumption rate in a dose-dependent manner (Fig. 2E), showing linear correlation with the RCR for ADP (Fig. 2E). However, Ppc-1 does not serve as a respiration substrate itself, and its oxygen consumption kinetics show that it does not affect ATP synthase (Fig. 2F). These results suggest that Ppc-1 is a novel mitochondrial uncoupler.




Figure 6. Detection of Ppc-1 in cells and tissues. (A) RPE cells were treated with 10 μ M Ppc-1 for 12 h, and then lipid fraction was prepared from cells, and analyzed on HPLC-MS/MS system B. The positions of Ppc-1 ([M+H]⁺ = 356.19 m/z) and its derivatives are indicated. (B) Sera and tissues were collected from mice at the time points indicated after a single shot of Ppc-1 (0.8 mg/kg), and the lipid fraction was prepared and analyzed similarly. The peak areas were calculated, and the results are expressed as mean and SD.

doi:10.1371/journal.pone.0117088.g006

Ppc-1 induces weight loss

We evaluated the effects of Ppc-1 on weight and fat content by administering it directly to mice. We found that Ppc-1 significantly suppresses weight gain in mice. It is noteworthy that this effect of Ppc-1 is observed at 21°C, a typical vivarium temperature that is below thermoneutrality [31]. Ppc-1 also leads to higher serum amounts of fatty acids and triacylglycerols, with lower percentages of subcutaneous and visceral fat compared to control mice. However, no other serum parameters show significant differences between Ppc-1-treated and control mice. The serum levels of ALT, BUN and adiponectin are all normal, and we observed no abnormalities in liver and kidney tissues. We initially guessed that Ppc-1 might induce adverse effects such as growth inhibition, since Ppc-1 inhibited growth of cultured cells [19] (S3 Fig.). However, we found that Ppc-1 concentrations were very low in serum and in all tissues tested




Figure 7. Release of fatty acids from 3T3-L1 cells by Ppc-1. Lipid extracts in the culture supernatants of differentiated 3T3-L1 cells treated with 10 μ M Ppc-1 for 48 h or 72 h were analyzed by HPLC-MS/MS system. A. A representative analysis of total fatty acids (A), and fatty acid components (B) (72 h only) is shown. The peak areas were calculated, and results are expressed as mean and SD. ** $p < 0.01$.

doi:10.1371/journal.pone.0117088.g007

in mice that were administered the compound (Fig. 7). Furthermore, no ectopic fat was detected in Ppc-1-treated mice. Thus, we speculate that Ppc-1 functions as an uncoupler in adipocytes and other cells in Ppc-1-treated animals, reducing the efficiency of mitochondrial ATP production. No excess heat production was observed, possibly because the Ppc-1 absorption and uncoupling reaction proceeded slowly over the course of serial administration. Low concentrations of Ppc-1 in the liver and kidney suggest a rapid degradation or modification by metabolic pathways. On the other hand, it is interesting that Ppc-1 stimulates adipocyte-like 3T3-L1 cells to release fatty acids into the culture medium (Fig. 7). Although the mechanism underlying the fatty acid release is under investigation, these findings are likely to be closely related to the increased serum fatty acids observed in mice treated with Ppc-1.

Obesity is a complex multifactorial disease and is a risk factor for other chronic diseases, including type 2 diabetes and cardiovascular disease. Understanding the metabolic factors and mitochondria dysfunction associated with obesity and weight loss success is important for developing appropriate prevention and treatment strategies [32]. Drugs for weight loss have been sought by many studies. DNP is reported to cause rapid weight loss, but leads to an unacceptably high number of adverse side effects [33–35]. Amphetamine derivatives have been used as appetite suppressants, but induce side effects that include valvular heart disease and lung hypertension. A new centrally active appetite suppressant and a lipase inhibitor are currently

being used as diet pills, but have equally unpleasant side effects. A small molecule, AdipoRon, which binds to adiponectin receptors, is currently being developed for the treatment of type 2 diabetes and obesity [36]. Adiponectin is an anti-diabetic and anti-atherogenic adipokine that binds to adiponectin receptors AdipoR1 and AdipoR2, and exerts anti-diabetic effects *via* the activation of the AMPK and PPAR- α pathways. Since AdipoRon is orally active, a therapeutic approach using AdipoR agonists is promising for the treatment of diabetes and other obesity-related diseases. Alternatively, it has been reported that DNP-methyl ester safely decreases hypertriglyceridemia and insulin resistance without systemic toxicity in rats fed a high fat diet [37]. DNP-methyl ester is a liver-targeted derivative of DNP, demonstrating the potential feasibility of dissociating the toxicity of DNP from its efficacy by altering its pharmacokinetics. DNP-methyl ester is a promising new drug candidate for the prevention of nonalcoholic fatty acid liver and type 2 diabetes. In contrast to these compounds, the mechanism of action for Ppc-1 is obscure, and the effective dose range that induces significant weight loss is quite narrow; weight loss occurs at an intermediate dose of 0.8 mg/week/kg (group B mice), with less weight loss at higher doses of 4 mg/week/kg and 10 mg/week/kg, and no weight loss at a lower dose (0.16 mg/week/kg). Thus, it will be necessary to evaluate the effects of Ppc-1 in mice fed high-fat diets and other fatty liver models, as well as optimize the form of administration including oral and percutaneous routes.

Action mechanism of Ppc-1

The molecular targets of Ppc-1 and its uncoupling mechanisms remain under investigation. Preliminary results indicate that uncoupling by Ppc-1 does not depend on the uncoupling proteins, UCP1 and UCP2. Furthermore, the uncoupling effect of Ppc-1 must be independent of ATP production, including respiratory complex V, because there is no inhibitory effect on 'state 3'. We considered whether Ppc-1 stimulates the formation of the permeability transition pore (PTP) [38], resulting in proton inflow to the matrix and uncoupling. The PTP modulates increased mitochondrial inner membrane permeability to ions and solutes with molecular masses up to about 1,500 Da, which causes matrix swelling. Thus, the PTP is central to vital mitochondrial functions and can play a lethal role under many pathophysiological conditions. However, Ppc-1 does not stimulate mitochondrial swelling nor apoptosis; therefore, the PTP is not likely to be the target of Ppc-1. Further studies are needed to address the molecular mechanism of Ppc-1.

Conclusions

Ppc-1 derived from slime molds was isolated from a library of small molecules as a novel agent which stimulates oxygen consumption in mitochondria with no inhibitory effects on ATP production. Direct administration of Ppc-1 to mice suppresses weight gain without abnormal changes in liver and kidney tissues, and no detectable tumor formation. The levels of free fatty acids are significantly elevated in the serum of mice treated with Ppc-1, while body fat content is relatively low. Ppc-1 stimulates cultured adipocytes to release fatty acids into the medium, which might be related to the rise in fatty acid levels in the serum of Ppc-1-treated mice. Our results suggest that Ppc-1 is a unique uncoupling agent applicable as a molecular probe for mitochondria research, and likely to aid the development of new drugs for the treatment of obesity.

Methods

Animal experiments

All experiments were carried out with the approval of the Fukushima Medical University Animal Studies Committee. Female ICR mice were obtained from CLEA Japan (Tokyo, Japan),

and housed at 21°C with a 12:12 h light:dark cycle with free access to water and a commercial diet. Ppc-1 dissolved in dimethyl sulfoxide was diluted with PBS and injected once a week into the peritoneal cavity for 8 weeks. Injections started when body weight was over 25 g, and doses of Ppc-1 varied between 0 and 10 mg/kg. Body weight and food intake were measured every 2 days just prior to lights out. Serum was collected from each animal on day 5 after the final administration of Ppc-1, and serum components including the albumin:globulin ratio, triglycerides, free fatty acids, creatinine, γ -GTP, aspartate aminotransferase (AST) and urea-N (BUN) were measured. Three mice were selected at random from each group, and euthanized for body fat analysis and tissue examination. Body fat was analyzed by LaTheta LCT-200 CT (Hitachi-Aloka, Tokyo). Livers and kidneys were fixed in paraformaldehyde and 6- μ m-sections were prepared. For detection of Ppc-1 in various tissues, liver, kidney, visceral fat, and serum were collected from mice treated with 0.8 mg/kg Ppc-1 for various time periods. Body surface temperature was monitored with an infrared thermometer.

Chemicals

The compounds screened in this study are listed in Fig. 1A; all were derived from *Dictyostelium* cellular slime molds. These compounds were synthesized and purified as described [8,9,20–24], and the structures and purities were confirmed by ^1H and ^{13}C NMR spectroscopy and high resolution MS. The purities of all compounds were greater than 98%. Oligomycin A, palmitoyl-L-carnitine, and fat-free bovine serum albumin (BSA) were purchased from Sigma-Aldrich.

Preparation of mitochondria-enriched fractions

Mitochondria were isolated from mouse liver (ICR; 7–10 week old females) by differential centrifugation as described previously [28]. Cells were homogenized in a Potter glass homogenizer with H-Buffer (250 mM sucrose, 10 mM Tris-HCl, pH 7.4), and centrifuged at 800xg for 1 min at 4°C. The supernatant was centrifuged at 6,000xg for 5 min, and the resulting pellet, the crude mitochondrial fraction, was suspended in H-Buffer. The suspension was layered over a discontinuous sucrose gradient consisting of 1.0 M and 1.5 M sucrose in Tris buffer (10 mM Tris-HCl, pH 7.4), and centrifuged for 20 min at 26,000xg at 4°C. The interface was collected in Tris buffer and centrifuged at 6,000xg for 5 min. The resulting pellets were suspended in Tris buffer (3 mg protein/ml) and used for experiments after confirming the presence of the mitochondrial marker cytochrome c.

Oxygen consumption analysis

Mitochondrial oxygen consumption was measured using a Clark-type oxygen electrode (Strathkelvin Instruments, North Lanarkshire, Scotland) as described [28–30,39]. Prior to the assay, an aliquot of mitochondrial-enriched fraction (0.1 mg protein) was incubated at 30°C in oxygen measurement buffer (225 mM mannitol, 75 mM sucrose, 10 mM KCl, 0.1 mM EDTA, 3 mM phosphate, 100 μ M palmitoyl-L-carnitine, and 20 mM Tris-HCl, pH 7.4) for 0.5 min. A test compound was added to the mixture at time = 0. Recording of oxygen level was started 30 s after the addition of the compound, and then an aliquot of ADP was added to a final concentration of 400 μ M to induce 'state 3'. Oligomycin A and KCN were used as respiratory system inhibitors.

Cell culture

RPE cells were grown and maintained at 37°C under 5% CO_2 and 95% air in Dulbecco's modified Eagle's medium-low glucose (DMEM-low glucose, 5 mM glucose) supplemented with 10%

(v/v) heat-inactivated fetal bovine serum (FBS) in a humidified incubator. 3T3-L1 fibroblasts were maintained in DMEM-high glucose (25 mM) containing 10% FBS. Post-confluent cells were differentiated into adipocytes by culturing for 3 days in medium containing 1 μ g/ml insulin, 0.1 μ g/ml dexamethasone, and 112 μ g/ml isobutylmethylxanthine [40,41]. Then the differentiation medium was replaced with DMEM, and the cells were cultured further for 10–14 days with medium changes every 4 days. Approximately 90% of cells showed the presence of large lipid droplets. Prior to the addition of Ppc-1, cells were maintained in a serum-free, BSA-containing (2%, w/v) medium for 1 day [41].

Mass spectrometry

Fatty acids and Ppc-1 in sera, tissues, and cells were analyzed by C18 reverse-phase liquid chromatography coupled to high resolution MS as reported [42,43]. Briefly, lipids and organic compounds were extracted by the method of Bligh and Dyer [44]. The free fatty acid compositions were analyzed on high performance liquid chromatography (HPLC)-MS/MS system A (HPLC, UltiMate 3000; MS, TSQ Vantage, Thermo Scientific) controlled by XCALIBUR (version 2.2, Thermo Scientific). To assess relative concentrations, the peak areas obtained from the MRM chromatogram for each fatty acid were normalized to the peak area of C17:0 margaric acid added as an internal standard. For the detection of Ppc-1, the extracts were analyzed on HPLC-MS/MS system B (HPLC, UltiMate 3000; MS, Orbitrap, Thermo Scientific) controlled by XCALIBUR. The *m/z* list of Ppc-1 was obtained from Metworks (version 1.3, Rococo Co., Ltd, Osaka, Japan) software for the prediction of *m/z* of drugs and their derivatives. The peak values of Ppc-1 were calculated from mass spectra obtained by positive acquisition polarity in a mass spectrometer.

Statistical analysis

Data were assessed using ANOVA with Tukey-Kramer post-hoc comparisons, and values are expressed as mean and SD unless otherwise indicated. The weights and fat levels of mice in the control and treated groups were evaluated by the Mann-Whitney test. Two-tailed values with $p < 0.05$ were considered statistically significant.

Supporting Information

S1 Fig. Weight changes following Ppc-1 treatment. Ppc-1 was injected once a week into the peritoneal cavity for 8 weeks. Doses of Ppc-1 for group A and group D were 0.16 (A) and 10 (B) mg/week/kg. No significant weight loss was observed in these groups.
(TIF)

S2 Fig. Body fat levels were examined using a CT scanner; representative cross section images of control and group B mice are shown ($n = 3$, each group). The position analyzed is denoted by an arrow in Fig. 4B.
(TIF)

S3 Fig. Effect of Ppc-1 on cell growth and ATP metabolism. (A) Cells were treated with the indicated doses of Ppc-1 for 48 h, and cell numbers were determined by counting. (B) ATP synthase activity was determined using the mitochondrial-enriched fraction. After 5 min incubation in oxygen measurement buffer containing the indicated amounts of Ppc-1, ADP was added to the reaction mixture to induce 'state 3'. ATP contents were measured 1 min before and 1 min after the addition of ADP, and the ATP synthesis rates were calculated from the difference in the two values. (C) ATPase activity was determined using mitochondria-enriched fraction, and the enzyme activity was measured as reported [45].
(TIF)

S1 Table. Body Fat and Profiling of Serum Factors. (DOCX)

Acknowledgments

The authors thank Drs. N Ahn and M Ohto for critical reading of the manuscript.

Author Contributions


Conceived and designed the experiments: YH MH HK YO. Performed the experiments: TS HK MO YH. Analyzed the data: TS MO. Contributed reagents/materials/analysis tools: HK YO. Wrote the paper: YH HK MO MH.

References

1. Singer TP (1979) Mitochondrial electron-transport inhibitors. *Methods Enzymol* 55: 454–462. PMID: [223000](#)
2. Heytler PG (1979) Uncouplers of oxidative phosphorylation. *Methods Enzymol* 55: 462–472. PMID: [156853](#)
3. Cooper CE, Brown GC (2008) The inhibition of mitochondrial cytochrome oxidase by the gases carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide: chemical mechanism and physiological significance. *J Bioenerg Biomembr* 40: 533–539. doi: [10.1007/s10863-008-9166-6](#) PMID: [18839291](#)
4. Horgan DJ, Singer TP, Casida JE (1968) Studies on the respiratory chain-linked reduced nicotinamide adenine dinucleotide dehydrogenase. 13. Binding sites of rotenone, piericidin A, and amytal in the respiratory chain. *J Biol Chem* 243: 834–843. PMID: [4295606](#)
5. Potter VR, Reif AE (1952) Inhibition of an electron transport component by antimycin A. *J Biol Chem* 194: 287–297. PMID: [14927618](#)
6. Loomis WF, Lipmann F (1948) Reversible inhibition of the coupling between phosphorylation and oxidation. *J Biol Chem* 173: 807–808. PMID: [18910739](#)
7. Heytler PG (1963) uncoupling of oxidative phosphorylation by carbonyl cyanide phenylhydrazones. I. Some characteristics of m-Cl-CCP action on mitochondria and chloroplasts. *Biochemistry* 2: 357–361. PMID: [13954345](#)
8. Morris HR, Taylor GW, Masento MS, Jermyn KA, Kay RR (1987) Chemical structure of the morphogen differentiation inducing factor from *Dictyostelium discoideum*. *Nature* 328: 811–814. PMID: [3627228](#)
9. Morris HR, Masento MS, Taylor GW, Jermyn KA, Kay RR (1988) Sstructure elucidation of two differentiation inducing factors (DIF-2 and DIF-3) from the cellular slime mold *Dictyostelium discoideum*. *Biochem J* 249: 903–906. PMID: [3355503](#)
10. Asahi K, Sakurai A, Takahashi N, Kubohara Y, Okamoto K, et al. (1995) DIF-1, morphogen of *Dictyostelium discoideum*, induces the erythroid differentiation in murine and human leukemia cells. *Biochem Biophys Res Commun* 208: 1036–1039. PMID: [7702602](#)
11. Kubohara Y, Saito Y, Tatemoto K (1995) Differentiation-inducing factor of *D. discoideum* raises intracellular calcium concentration and suppresses cell growth in rat pancreatic AR42J cells. *FEBS Lett* 359: 119–122. PMID: [7867781](#)
12. Kubohara Y (1997) DIF-1, putative morphogen of *D. discoideum*, suppresses cell growth and promotes retinoic acid-induced cell differentiation in HL-60. *Biochem Biophys Res Commun* 236: 418–422. PMID: [9240452](#)
13. Kubohara Y (1999) Effects of differentiation-inducing factors (DIFs) of *Dictyostelium discoideum* on human leukemia K562 cells: DIF-3 is the most potent anti-leukemic agent. *Eur J Pharmacol* 381: 57–62. PMID: [10528134](#)
14. Kanai M, Konda Y, Nakajima T, Izumi Y, Kanda N, et al. (2003) Differentiation-inducing factor-1 (DIF-1) inhibits STAT3 activity involved in gastric cancer cell proliferation via MEK-ERK dependent pathway. *Oncogene* 22: 548–554. PMID: [12555068](#)
15. Takahashi-Yanaga F, Taba Y, Miwa Y, Kubohara Y, Watanabe Y, et al. (2003) *Dictyostelium* differentiation-inducing factor-3 activates glycogen synthase kinase-3 β and degrades cyclin D1 in mammalian cells. *J Biol Chem* 278: 9663–9670. PMID: [12522140](#)
16. Takahashi-Yanaga F, Mori J, Matsuzaki E, Watanabe Y, Hirata M, et al. (2006) Involvement of GSK-3 β and DYRK1B in differentiation-inducing factor-3-induced phosphorylation of cyclin D1 in HeLa cells. *J Biol Chem* 281: 38489–38497. PMID: [17046823](#)

17. Kubohara Y, Kikuchi H, Matsuo Y, Oshima Y, Homma Y (2013) Mitochondria are the target organelle of differentiation-inducing factor-3, an anti-tumor agent isolated from *Dictyostelium Discoideum*. PLoS One 8: e72118. doi: [10.1371/journal.pone.0072118](https://doi.org/10.1371/journal.pone.0072118) PMID: [23977224](https://pubmed.ncbi.nlm.nih.gov/23977224/)
18. Kubohara Y, Kikuchi H, Matsuo Y, Oshima Y, Homma Y (2014) Properties of a non-bioactive fluorescent derivative of differentiation-inducing factor-3, an anti-tumor agent found in *Dictyostelium discoideum*. Biol Open 3: 289–296. doi: [10.1242/bio.20146585](https://doi.org/10.1242/bio.20146585) PMID: [24682009](https://pubmed.ncbi.nlm.nih.gov/24682009/)
19. Kikuchi H, Ishiko S, Nakamura K, Kubohara Y, Oshima Y (2010) Novel prenylated and geranylated aromatic compounds isolated from *Polysphondylium* cellular slime molds. Tetrahedron 66: 6000–6007.
20. Takaya Y, Kikuchi H, Terui Y, Komiya J, Furukawa K-I, et al. (2000) Novel acyl α -pyronoids, dictyopyrone A, B, and C, from *Dictyostelium* cellular slime molds. J Org Chem 65: 985–989. PMID: [10814044](https://pubmed.ncbi.nlm.nih.gov/10814044/)
21. Kikuchi H, Nakamura K, Kubohara Y, Gokan N, Hosaka K, et al. (2007) Dihydrodictyopyrone A and C: new members of dictyopyrone family isolated from *Dictyostelium* cellular slime molds. Tetrahedron Lett 48: 5905–5909.
22. Kikuchi H, Saito Y, Kojima J, Takaya Y, Honma S, et al. (2001) Furanodictine A and B: amino sugar analogues produced by cellular slime mold *Dictyostelium discoideum* showing neuronal differentiation activity. J Org Chem 66: 6982–6987. PMID: [11597217](https://pubmed.ncbi.nlm.nih.gov/11597217/)
23. Kikuchi H, Komiya J, Saito Y, Sekiya J-I, Honma S, et al. (2002) The isolation and synthesis of two novel N-acetyl glucosamine derivatives from *Dictyostelium* slime molds which exhibit neurite outgrowth activity. Tetrahedron Lett 43: 1477–1480.
24. Kikuchi H, Saito Y, Sekiya J, Okano Y, Saito M, et al. (2005) Isolation and synthesis of a new aromatic compound, brefelamide, from *Dictyostelium* slime molds and its inhibitory effect on the proliferation of astrocytoma cells. J Org Chem 70: 8854–8858. PMID: [16238318](https://pubmed.ncbi.nlm.nih.gov/16238318/)
25. Matsuno-Yagi A, Hatefi Y (1986) Kinetic modalities of ATP synthesis. J Biol Chem 261: 14031–14038. PMID: [2945814](https://pubmed.ncbi.nlm.nih.gov/2945814/)
26. Hekman C, Matsuno-Yagi A, Hatefi Y (1988) Energy-induced modulation of the kinetics of oxidative phosphorylation and reverse electron transfer. Biochemistry 27: 7559–7565. PMID: [2905168](https://pubmed.ncbi.nlm.nih.gov/2905168/)
27. Guzy RD, Sharma B, Bell E, Chandel NS, Schumacker PT (2008) Loss of the SdhB, but not the SdhA, subunit of complex II triggers reactive oxygen species-dependent hypoxia-inducible factor activation and tumorigenesis. Mol Cell Biol 28: 718–731. PMID: [17967865](https://pubmed.ncbi.nlm.nih.gov/17967865/)
28. Kabuyama Y, Suzuki T, Nakazawa N, Yamaki J, Homma MK, et al. (2010) Dysregulation of very long chain acyl-CoA dehydrogenase coupled with lipid peroxidation. Am J Physiol Cell Physiol 298: C107–C113. doi: [10.1152/ajpcell.00231.2009](https://doi.org/10.1152/ajpcell.00231.2009) PMID: [19889959](https://pubmed.ncbi.nlm.nih.gov/19889959/)
29. Ogura M, Yamaki J, Homma MK, Homma Y (2012) Mitochondrial c-Src regulates cell survival through phosphorylation of respiratory chain components. Biochem J 447: 281–289. PMID: [22823520](https://pubmed.ncbi.nlm.nih.gov/22823520/)
30. Ogura M, Yamaki J, Homma MK, Homma Y (2014) Phosphorylation of flotillin-1 by mitochondrial c-Src is required to prevent the production of reactive oxygen species. FEBS Lett 588: 2837–2843. doi: [10.1016/j.febslet.2014.06.044](https://doi.org/10.1016/j.febslet.2014.06.044) PMID: [24983503](https://pubmed.ncbi.nlm.nih.gov/24983503/)
31. Goldgof M, Xiao C, Chanturiya T, Jou W, Gavrilova O, et al. (2014) The chemical uncoupler 2,4-dinitrophenol (DNP) protects against diet-induced obesity and improves energy homeostasis in mice at thermoneutrality. J Biol Chem 289: 19341–19350. doi: [10.1074/jbc.M114.568204](https://doi.org/10.1074/jbc.M114.568204) PMID: [24872412](https://pubmed.ncbi.nlm.nih.gov/24872412/)
32. Thrush AB, Dent R, McPherson R, Harper ME (2013) Implications of mitochondrial uncoupling in skeletal muscle in the development and treatment of obesity. FEBS J 280: 5015–5029. doi: [10.1111/febs.12399](https://doi.org/10.1111/febs.12399) PMID: [23786211](https://pubmed.ncbi.nlm.nih.gov/23786211/)
33. Grundlingh J, Dargan PI, El-Zanfaly M, Wood DM (2011) 2,4-Dinitrophenol (DNP): a weight loss agent with significant acute toxicity and risk of death. J Med Toxicol 7: 205–212. doi: [10.1007/s13181-011-0162-6](https://doi.org/10.1007/s13181-011-0162-6) PMID: [21739343](https://pubmed.ncbi.nlm.nih.gov/21739343/)
34. Poole FE, Haining RB (1934) Sudden death from dinitrophenol poisoning. JAMA 102: 1141–1147.
35. Tainter ML, Wood DA (1934) A case of fatal dinitrophenol poisoning. JAMA 102: 1147–1149.
36. Okada-Iwabu M, Yamauchi T, Iwabu M, Honma T, Hamagami K, et al. (2013) A small-molecule AdipoR against for type 2 diabetes and short life in obesity. Nature 503: 493–499. doi: [10.1038/nature12656](https://doi.org/10.1038/nature12656) PMID: [24172895](https://pubmed.ncbi.nlm.nih.gov/24172895/)
37. Perry RJ, Kim T, Zhang X-M, Lee H-Y, Pesta D, et al. (2013) Reversal of hypertriglyceridemia, fatty liver diseases, and insulin resistance by a liver-targeted mitochondrial uncoupler. Cell Metabolism 18: 740–748. doi: [10.1016/j.cmet.2013.10.004](https://doi.org/10.1016/j.cmet.2013.10.004) PMID: [24206666](https://pubmed.ncbi.nlm.nih.gov/24206666/)
38. Bernardi P (2013) The mitochondrial permeability transition pore: a mystery solved? Front Physiol. 4: 95. doi: [10.3389/fphys.2013.00095](https://doi.org/10.3389/fphys.2013.00095) PMID: [23675351](https://pubmed.ncbi.nlm.nih.gov/23675351/)


39. Katou Y, Endo N, Suzuki T, Yu J, Kikuchi H, et al. (2014) Matarhizin A suppresses cell proliferation by inhibiting cytochrome c oxidase activity. *Life Sci* 103: 1–7. doi: [10.1016/j.lfs.2014.03.023](https://doi.org/10.1016/j.lfs.2014.03.023) PMID: [24699005](https://pubmed.ncbi.nlm.nih.gov/24699005/)
40. Worrall DS, Olefsky JM (2002) The effects of intracellular calcium depletion on insulin signaling in 3T3-L1 adipocytes. *Mol Endocrinol* 16: 378–389. PMID: [11818508](https://pubmed.ncbi.nlm.nih.gov/11818508/)
41. Kudoh A, Satoh H, Hirai H, Watanabe T (2011) Pioglitazone upregulates adiponectin receptor 2 in 3T3-L1 adipocytes. *Life Sci* 88: 1055–1062. doi: [10.1016/j.lfs.2011.04.001](https://doi.org/10.1016/j.lfs.2011.04.001) PMID: [21514306](https://pubmed.ncbi.nlm.nih.gov/21514306/)
42. Yamada T, Uchikata T, Sakamoto S, Yokoi Y, Fukusaki E, et al. (2013) Development of a lipid profiling system using reverse-phase liquid chromatography coupled to high-resolution mass spectrometry with rapid polarity switching and an automated lipid identification software. *J Chromatogr A* 1292: 211–218. doi: [10.1016/j.chroma.2013.01.078](https://doi.org/10.1016/j.chroma.2013.01.078) PMID: [23411146](https://pubmed.ncbi.nlm.nih.gov/23411146/)
43. Ohira H, Fujioka Y, Katagiri C, Mamoto R, Aoyama-Ishikawa M, et al. (2013) Butyrate attenuates inflammation and lipolysis generated by the interaction of adipocytes and macrophages. *J Atheroscler Thromb* 20: 425–442. PMID: [23470566](https://pubmed.ncbi.nlm.nih.gov/23470566/)
44. Bligh EG, Dyer WJ (1959) A rapid method for total lipid extraction and purification. *Can J Biochem Physiol* 37: 911–917. PMID: [13671378](https://pubmed.ncbi.nlm.nih.gov/13671378/)
45. Rigoulet M, Devin A, Averet N, Vandais B, Guerin B (1996) Mechanisms of inhibition and uncoupling of respiration in isolated rat liver mitochondria by the general anesthetic 2,6-diisopropylphenol. *Eur J Biochem* 241: 280–285. PMID: [8898917](https://pubmed.ncbi.nlm.nih.gov/8898917/)




S1 Fig. Weight changes following Ppc-1 treatment.

Ppc-1 was injected once a week into the peritoneal cavity for 8 weeks. Doses of Ppc-1 for group A and group D were 0.16 (A) and 10 (B) mg/week/kg. No significant weight loss was observed in these groups.

doi:10.1371/journal.pone.0117088.s001



S2 Fig. Body fat levels were examined using a CT scanner; representative cross section images of control and group B mice are shown (n = 3, each group). The position analyzed is denoted by an arrow in Fig. 4B.
doi:10.1371/journal.pone.0117088.s002



S3 Fig. Effect of Ppc-1 on cell growth and ATP metabolism.

(A) Cells were treated with the indicated doses of Ppc-1 for 48 h, and cell numbers were determined by counting. (B) ATP synthase activity was determined using the mitochondrial-enriched fraction. After 5 min incubation in oxygen measurement buffer containing the indicated amounts of Ppc-1, ADP was added to the reaction mixture to induce 'state 3'. ATP contents were measured 1 min before and 1 min after the addition of ADP, and the ATP synthesis rates were calculated from the difference in the two values. (C) ATPase activity was determined using mitochondria-enriched fraction, and the enzyme activity was measured as reported [45].

doi:10.1371/journal.pone.0117088.s003

S1 Table. Body Fat and Profiling of Serum Factors

	Percent body fat (%)		Total cholesterol (mg/l)		Phospholipid (mg/l)		Albumin / Globulin		Creatinine (mg/l)		γ-GTP (IU/l)
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean
Control	15.1	2.9	778	121	1,302	164	1.88	0.10	1.50	0.61	<3
Group A	15.1	1.8	820	123	1,407	179	1.93	0.12	1.10	0.26	<3
Group B	11.3	4.5	708	81	1,290	168	1.70	0.18	1.68	0.48	<3
Group C	14.5	1.1	788	158	1,340	144	1.88	0.21	1.58	0.38	<3
Group D	17.9	2.6	740	158	1,305	288	2.03	0.22	1.53	0.34	<3